



**ST VINCENT'S
PRIVATE HOSPITAL**
SYDNEY

CONDUCTED BY THE SISTERS OF CHARITY OF AUSTRALIA

Dr Rohan M Gett, FRACS

Colorectal Surgeon

St. Vincent's Clinic

438 Victoria St, Darlinghurst NSW 2010

T: 8382 6675

Home Care Advice Post Bowel Resection

DO's

- ❖ Make an appointment with Dr Gett for 2-4 weeks after your surgery.
- ❖ Continue to walk short distances, including stairs, regularly.
- ❖ If you still have waterproof dressing/s leave them on for 5 days, then remove them and leave your wound/s exposed, ensuring you dry them thoroughly after showering.
- ❖ If you still have drains/s empty and record the output at least daily or when full. Ensure the drain is on suction at all times. (nursing staff will instruct you on this)
- ❖ Ask Dr Gett's secretary for a medical certificate if required.
- ❖ Organize at least 2-4 weeks away from work after your surgery.
- ❖ Take your pain medication as directed, if needed. Be aware that it may make you drowsy and possibly cause constipation.
- ❖ Eat a varied well balanced diet with plenty of water, some people find eating small frequent meals more beneficial than 3 large meals per day.



**ST VINCENT'S
PRIVATE HOSPITAL**
SYDNEY

CONDUCTED BY THE SISTERS OF CHARITY OF AUSTRALIA

Do's Continued

- ❖ Expect to feel tired and fatigue easily, this is normal and will improve over the next 1-2 weeks.
- ❖ Wear your TED stockings for at least the first week at home.
- ❖ Make an appointment with your GP within the first 2 weeks of being home to reassess your regular medications.
- ❖ Expect your bowels to be unpredictable for at least 1 month after your surgery.

Don'ts

- ❖ Do not drive for at least 2 whole weeks after your surgery.
- ❖ Do not lift more than 10kgs for 2-4 weeks post your surgery.
- ❖ Do not have a bath; shower only until confirmed by Dr Gett.
- ❖ Do not return to strenuous exercise such as the gym, golf, running, swimming etc for at least 4 weeks post your surgery.



**ST VINCENT'S
PRIVATE HOSPITAL**
SYDNEY

CONDUCTED BY THE SISTERS OF CHARITY OF AUSTRALIA

**Please Call Your Doctor If You Notice Any Of The
Following:**

- ❖ **Increased swelling, redness or tenderness at your wound/s site.**
- ❖ **Any discharge from your wound/s.**
- ❖ **Increased pain.**
- ❖ **Fever or chills**
- ❖ **Unrelieved constipation or difficulty passing urine.**
- ❖ **Persistent nausea, vomiting &/or loss of appetite.**
- ❖ **Unpredictable and loose stools are to be expected for a short period. If loose frequent stools persist please call your doctor.**